



Evidence for Degenerate Tetraploidy in Bdelloid Rotifers

Citation

Welch, David B. Mark, Jessica L. Mark Welch, and Matthew Meselson. 2008. Evidence for degenerate tetraploidy in bdelloid rotifers. *Proceedings of the National Academy of Sciences USA* 105(13): 5145-5149

Published Version

<http://dx.doi.org/10.1073/pnas.0800972105>

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:3109360>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Evidence for degenerate tetraploidy in bdelloid rotifers

David B. Mark Welch*, Jessica L. Mark Welch*, and Matthew Meselson*^{††}

*Josephine Bay Paul Center for Comparative Molecular Biology and Evolution, Marine Biological Laboratory, Woods Hole, MA 02543; and [†]Department of Molecular and Cellular Biology, Harvard University, Cambridge, MA 02138

Contributed by Matthew Meselson, January 31, 2008 (sent for review November 9, 2007)

Rotifers of class Bdelloidea have evolved for millions of years apparently without sexual reproduction. We have sequenced 45- to 70-kb regions surrounding the four copies of the *hsp82* gene of the bdelloid rotifer *Philodina roseola*, each of which is on a separate chromosome. The four regions comprise two colinear gene-rich pairs with gene content, order, and orientation conserved within each pair. Only a minority of genes are common to both pairs, also in the same orientation and order, but separated by gene-rich segments present in only one or the other pair. The pattern is consistent with degenerate tetraploidy with numerous segmental deletions, some in one pair of colinear chromosomes and some in the other. Divergence in 1,000-bp windows varies along an alignment of a colinear pair, from zero to as much as 20% in a pattern consistent with gene conversion associated with recombinational repair of DNA double-strand breaks. Although pairs of colinear chromosomes are a characteristic of sexually reproducing diploids and polyploids, a quite different explanation for their presence in bdelloids is suggested by the recent finding that bdelloid rotifers can recover and resume reproduction after suffering hundreds of radiation-induced DNA double-strand breaks per oocyte nucleus. Because bdelloid primary oocytes are in G₁ and therefore lack sister chromatids, we propose that bdelloid colinear chromosome pairs are maintained as templates for the repair of DNA double-strand breaks caused by the frequent desiccation and rehydration characteristic of bdelloid habitats.

asexual reproduction | genome structure | evolution of sex | gene conversion | anhydrobiosis

There are numerous hypotheses and theoretical models but no general agreement about what selective factors and underlying mechanisms are responsible for the nearly universal occurrence of sexual reproduction among animals and plants and the relatively early extinction of lineages that abandon it (1–3). Ancient taxa in which sexual reproduction is unknown have therefore attracted attention as systems whose study may shed light on what has allowed them to avoid extinction and successfully evolve. Of these, the group for which evidence of ancient asexuality is strongest is the class Bdelloidea of the phylum Rotifera (4).

Bdelloid rotifers are common invertebrates a few tenths of a millimeter long characteristically found in the water films of mosses and lichens, in temporary freshwater pools, and in other ephemeral aquatic habitats. They are able to thrive in such settings because they can survive desiccation at any stage of their life cycle by entering a metabolically quiescent state of anhydrobiosis (5). Distinguished by their ciliated head structure, bilateral ovaries, and jaw-like mastax, they have ganglia; muscles; digestive and secretory systems; photosensitive and tactile sensory organs; and structures for feeding, crawling, and swimming (6, 7). The class includes 461 described species classified in four families and 19 genera (8). The identification to the family level of bdelloid remains in 35- to 40-million-year-old amber, the high synonymous site difference between bdelloid species of different families, and the molecular clock estimate that bdelloids separated from their sister class, the facultatively sexual monogonont

rotifers, at least 100 million years ago all suggest that the class is of Mesozoic origin (9–13).

Despite much observation and study of bdelloid rotifers in the field and in laboratory culture since they were first observed >300 years ago, males, hermaphrodites, vestigial male structures, and meiosis have never been documented within the class (14–16). Eggs are produced from primary oocytes by two mitotic divisions, without chromosome pairing and without reduction in chromosome number (17, 18).

One of the genes that has been characterized most extensively in bdelloids is the heat-shock gene *hsp82* of *Philodina roseola*, of which there are four copies, each on a separate chromosome (19, 20). Here we describe the sequence of the 45- to 70-kb region surrounding each copy of *hsp82* in *P. roseola*. The four regions are found to consist of two colinear, gene-rich pairs. The two pairs have only a minority of genes in common, and synonymous divergence between these genes is consistently much higher than that within a pair. The genes common to all four sequenced regions are in the same order and orientation and are separated by segments present in only one pair or the other. The arrangement therefore suggests that the genome is that of a degenerate tetraploid. Our findings are discussed in relation to bdelloid genome structure and its evolution, the adaptation of bdelloids to ephemerally aquatic habitats, and their putative ancient asexuality.

Results

The 45- to 70-kb regions surrounding each of the four copies of *hsp82* in *P. roseola* are depicted in Fig. 1, and annotations and divergence values are listed in Table 1. It is seen that (i) the four contigs may be grouped in a quartet of two colinear pairs, with gene content, order, and orientation conserved within each pair. The two pairs are designated A and B, respectively, corresponding to the A and B pairs of *hsp82* genes we previously identified (19, 20). (ii) The average divergence between members of a pair in 1-kb sliding windows is $3.3 \pm 2.6\%$ and $5.3 \pm 3.8\%$ for the A and B pairs, respectively, and varies widely along their length: there are regions as long as 1 kb that differ by as much as 20% and tracts of identity or near-identity up to several kilobases in length. (iii) Of the 16 genes identified in the region bracketed by the two outside genes common to both pairs (*plasma membrane calcium ATPase* and *SAICAR synthetase-AIR carboxylase*), only two other genes are present on both pairs (*myosin light chain* and *hsp82*). The four common genes are in the same order and orientation on each of the four contigs and are separated by genes that are present in one pair but absent in the other. (iv) Coding sequences of identified genes make up $\approx 50\%$ of the sequenced regions. (v) Synonymous site

Author contributions: D.B.M.W. and J.L.M.W. designed research; D.B.M.W. and J.L.M.W. performed research; D.B.M.W. and M.M. analyzed data; and D.B.M.W. and M.M. wrote the paper.

The authors declare no conflict of interest.

Data deposition: The sequences reported in this paper have been deposited in the GenBank database (accession nos. EU432546–EU432549).

^{††}To whom correspondence should be addressed. E-mail: msm@wjh.harvard.edu.

© 2008 by The National Academy of Sciences of the USA




Fig. 1. The four contigs containing *hsp82* in *P. roseola*. Maps of the four contigs are shown between A and B; copies 1 and 2 belong to lineage A, and copies 3 and 4 belong to lineage B. Filled rectangles represent genes. Gene orientation is indicated by position above or below the line, and numbers refer to genes listed in Table 1. Genes present in all four contigs are connected by dashed lines; the alignment is centered on *hsp82* and is to scale with the graphs in A and B. The graphs show divergence along the alignment of the A pair (A) and B pair (B) in 100-bp (thin lines) and 1,000-bp (thick lines) sliding windows, with the average Ks of each gene pair shown as a horizontal thick line.

divergence (Ks) of genes within a pair displays a wide range of values, from zero to $\approx 20\%$. The nonweighted average Ks is 7.4%, more than three standard deviations greater than the synonymous and silent site diversity of 2.7% averaged over diverse regions in a large number of invertebrate species and greater than the average Ks recorded for any one species (21).

Additional findings, not depicted in the figure, are that (vi) introns are numerous and typically 50–60 bp long, with conserved GT-AG boundaries. An intron in copy 2 of *methionine-sulfoxide reductase B* is absent in copy 1 of the gene, and in *plasma membrane calcium ATPase* the A pair and the B pair each share an intron absent in the other pair. (vii) Except for a possible foldback element in one of the B contigs (22), there is no evidence of intact or relict transposons or pseudogenes in any of the four contigs. (viii) Within both colinear pairs there are numerous small indels, a few as large as 500 bp. (ix) The A sequences cannot be aligned with the B sequences except in the portions corresponding to the four genes that they share. (x) Ks between A and B copies of these four genes is 112%, 68%, 67%, and 122%, respectively. (xi) There are no tracts of identity or near identity between A and B sequences. (xii) There are no significant differences between contigs in overall or third position GC content or in codon bias. (xiii) No identified gene contains frameshifts or internal stop codons. (xiv) Ka/Ks ranges from 0.03 to 0.59, indicating the operation of purifying selection on amino acid sequence.

Discussion

Bdelloid Genome Structure. We find that the sequences around the four copies of *hsp82* in *P. roseola* occur as two colinear pairs, a

quartet in which the two pairs share a limited number of genes in the same order and orientation but with genes present in each pair that are absent from the other. That the quartet structure of the *hsp82* region of *P. roseola* is conserved in Bdelloidea is supported by the finding of an *hsp82*-containing quartet with a gene content and structure closely similar to that depicted in Fig. 1 in the bdelloid *Adineta vaga* (23), a species belonging to a family that separated from that of *P. roseola* ≈ 60 million years ago (12, 19, 23). Moreover, the existence of such quartet structures along entire chromosomes is suggested by the finding in *P. roseola* of a quartet organization of the regions surrounding a Hox gene and by the observation by FISH that the Hox and *hsp82* quartets appear to be far apart on the same four mid-size chromosomes (J.L.M.W., unpublished observations).

That bdelloid genomes are generally organized as quartets is also consistent with the karyotypes of *P. roseola* and *A. vaga*. The karyotype of *A. vaga* consists of 12 mid-size chromosomes, probably comprising three quartets, whereas that of *P. roseola* comprises 10 mid-size chromosomes, a single chromosome of approximately twice that length, and two dots of unequal size (18, 24). The large chromosome appears to be an isochromosome, because probes that hybridize to it do so at pairs of sites, each approximately equidistant from its center (ref. 25 and unpublished data). The 13 chromosomes of *P. roseola* may therefore consist of two quartets of mid-size chromosomes, with the two remaining mid-size chromosomes and the two arms of the isochromosome making up the third quartet. The two dots may be B chromosomes, because one or the other is often missing in karyotypes (J.L.M.W., unpublished observations).

Thus, the evidence suggests that bdelloids are degenerate

lineage contigs occurring anywhere in the region of their overlap is 3.0×10^{-6} , and that of the 400-bp tract in the B lineage is 2.8×10^{-7} . Inspection of the colinear pairs reveals additional long tracts that are nearly identical. Such tracts of identity and near-identity appear to have arisen recently, possibly associated with DSB repair, which may also produce tracts too short to be recognized as such. The correlation of divergence values between adjacent windows is very strong ($r \approx 0.7$) for windows up to ≈ 2 kb and becomes insignificant by about twice that length (data not shown). This pattern, seen over a wide range of divergence values, would be expected for gene conversion tracts usually of no more than a few kilobases occurring over millions of years. The contigs depicted in Fig. 1 exhibit no evidence of recent conversion between A and B lineages, and these are now probably too divergent to function as templates for interlineage homologous repair.

In addition to satisfying the requirement for templates for accurate DNA repair in G_1 , colinear chromosome pairs may be of additional benefit to bdelloid rotifers by allowing the accumulation of heterotic interactions. Because the average synonymous divergence we find for gene pairs on colinear chromosomes is $\approx 7\%$, such interactions could accumulate and persist for appreciable times between homogenization events. Also, the presence of colinear chromosome pairs and the occurrence of gene conversion may produce clones with combinations of mutations within a gene that are beneficial only in *cis* more rapidly than could be achieved if each chromosome were represented only once. Furthermore, given the requirement for colinear chromosome pairs, conversion and mitotic crossing-over in clones heterozygous for recessive beneficial mutations may hasten the appearance of better-adapted clones that otherwise could arise only by further mutation in the clone in which they occur, analogous to the effect of allelic segregation and syngamy in sexual species (38, 39).

Implications of Colinear Pairs for Bdelloid Asexuality. Our findings call for a reevaluation of the significance of the highly diverged pairs of genes we have found in individual bdelloid genomes (19). Previously, although not rejecting the possibility that bdelloids descend from an ancient polyploid ancestor and engage in some rare or elusive form of sexual reproduction, we thought it more plausible, consistent with other indications of bdelloid asexuality and the presumed rareness of polyploidy in animals (40), to interpret the highly diverged copies of *hsp82* we found in diverse bdelloid species as descendants of alleles that had ceased segregation and had not undergone homogenization by conversion or mitotic crossing-over since the separation of bdelloid families

and therefore as evidence for ancient asexuality (19). The present evidence that bdelloids are degenerate tetraploids suggests that the highly diverged gene pairs are the result of ancient whole-genome duplication, perhaps involving a hybridization between closely related species, rather than descendants of former alleles and raises the question of whether the colinear chromosomes are meiotic homologs. Although it is not entirely excluded that bdelloids engage in rare sexual reproduction, as explained above an entirely different explanation for the maintenance of colinear chromosome pairs follows from the discovery of the extraordinary resistance of bdelloid rotifers to ionizing radiation (32) and the implication that colinear chromosome pairs are required in order for bdelloids to repair DNA breakage associated with their ephemerally aquatic habitats.

Materials and Methods

For each of the four copies of *hsp82*, two cosmids with overlapping inserts were selected from a genomic library prepared from a single-egg culture of *P. roseola* as previously described (20). Each cosmid was digested to completion with two or three six-base recognition restriction endonucleases and subcloned in pBSKII (Stratagene). Plasmid DNA was extracted manually (41) or with a MiniPrep 24 (MacConnell Research) or RevPrep (Gene Machines) system. Sequences were obtained by using BigDye Terminator sequencing reagents and ABI automated capillary sequencers or the SequiTherm EXCEL II DNA sequencing kit (Epicentre) and a 4,000-IR² DNA scanner (LI-COR). Sequences of the overlapping cosmids were assembled into single contigs with six times average coverage using Sequencher (Gene Codes). Sequence gaps were filled by sequencing directly from 1 μ g of cosmid DNA using specifically designed primers. The identity of overlapping regions of cosmids from each contig indicated that the cosmid inserts are not chimeric. Contigs were aligned by using the EMBOSS program *stretcher*.

Gene regions were identified by BLASTX searches of *Drosophila melanogaster*, *Caenorhabditis elegans*, and the National Center for Biotechnology Information nonredundant databases followed by BLASTX and BLASTN searches of putative intergenic regions, by hidden Markov model searches of Pfam full-length and fragment domain models using HMMER (42), and by searches for ORFs. Exons and introns were mapped by comparison to homologous amino acid sequences using *genewise* (43). Where homology to annotated reference sequences was poor (particularly near exon boundaries), aligned contigs were scanned in all six reading frames for regions in which the ratio of nonsynonymous to synonymous difference is <1 .

Divergence at synonymous sites (Ks) and at nonsynonymous sites (Ka) was measured with the *diverge* program of the Wisconsin Package (Accelrys), which uses the method of Pamilo and Bianchi (44). Sliding window analysis of total divergence was done with DnaSP 4.0. The runs test of Stephens (37) was implemented by using Sterling's approximation for factorials.

ACKNOWLEDGMENTS. We thank the Eukaryotic Genetics Program of the U.S. National Science Foundation for support and the Bay Paul Center W. M. Keck Ecological and Evolutionary Genetics Sequencing Facility.

- Kondrashov AS (1993) Classification of hypotheses on the advantages of amphimixis. *J Hered* 84:372–387.
- Barton NH, Charlesworth B (1998) Why sex and recombination? *Science* 281:1986–1990.
- Otto SP, Lenormand T (2002) Resolving the paradox of sex and recombination. *Nat Rev Genet* 3:252–260.
- Normark BB, Judson OP, Moran NA (2003) Genomic signatures of ancient asexual lineages. *Biol J Linn Soc London* 79:69–84.
- Ricci C (1998) Anhydrobiotic capabilities of bdelloid rotifers. *Hydrobiologia* 387/388:321–326.
- Donner J (1965) *Ordnung Bdelloidea* (Akademie, Berlin).
- Hyman LH (1951) *The Invertebrates* (McGraw-Hill, New York).
- Segers H (2007) Annotated checklist of the rotifers (Phylum Rotifera), with notes on nomenclature, taxonomy and distribution. *Zootaxa* 1564:1–104.
- Poinar GO, Jr, Ricci C (1992) Bdelloid rotifers in Dominican amber: Evidence for parthenogenetic continuity. *Experientia* 48:408–410.
- Waggoner BM, Poinar GO, Jr (1993) Fossil habrotrichid rotifers in Dominican amber. *Experientia* 49:354–357.
- Mark Welch DB (2000) Evidence from a protein coding gene that acanthocephalans are rotifers. *Invertebr Biol* 111:17–23.
- Mark Welch DB, Meselson MS (2001) Rates of nucleotide substitution in sexual and asexually reproducing rotifers. *Proc Natl Acad Sci USA* 98:6720–6724.
- Mark Welch DB, Cummings MP, Hillis DM, Meselson M (2004) Divergent gene copies in the asexual class Bdelloidea (Rotifera) separated before the bdelloid radiation or within bdelloid families. *Proc Natl Acad Sci USA* 101:1622–1625.
- Leeuwenhoek A (1677) Letter of October 9, 1676 to the Royal Society. *Philos Trans R Soc London Ser B* 12:821–831.
- Hudson, CT, Gosse PH (1886) *The Rotifera or Wheel-Animalcules* (Longmans, Green, London).
- Ricci C (1987) The ecology of bdelloids: How to be successful. *Hydrobiologia* 147:117–127.
- Hsu WS (1956) Oogenesis in *Habrotricha tridens* (Milne). *Biol Bull* 111:364–374.
- Hsu WS (1956) Oogenesis in the Bdelloidea rotifer *Philodina roseola* Ehrenberg. *Cellule* 57:283–296.
- Mark Welch DB, Meselson M (2000) Evidence for the evolution of bdelloid rotifers without sexual reproduction or genetic exchange. *Science* 288:1211–1215.
- Mark Welch JL, Mark Welch DB, Meselson M (2004) Cytogenetic evidence for asexual evolution of bdelloid rotifers. *Proc Natl Acad Sci USA* 101:1618–1621.
- Lynch M (2006) The origins of eukaryotic gene structure. *Mol Biol Evol* 23:450–468.
- Arkhipova IR, Meselson M (2005) Diverse DNA transposons in rotifers of the class Bdelloidea. *Proc Natl Acad Sci USA* 102:11781–11786.
- Hur J (2006) Duplication and divergence of regions containing *hsp82* before the separation of two Bdelloid families, Adinetidae and Philodinidae. PhD dissertation (Harvard Univ, Cambridge, MA).
- Mark Welch JL, Meselson M (1998) Karyotypes of bdelloid rotifers from three families. *Hydrobiologia* 387/388:403–407.
- Mark Welch JL (2001) Cytological evidence for the absence of meiosis in bdelloid rotifers. PhD dissertation (Harvard Univ, Cambridge, MA).
- Wolfe KH (2001) Yesterday's polyploids and the mystery of diploidization. *Nat Rev Genet* 2:333–341.

27. Santos JL, et al. (2003) Partial diploidization of meiosis in autotetraploid *Arabidopsis thaliana*. *Genetics* 165:1533–1540.
28. Kellis M, Birren BW, Lander ES (2004) Proof and evolutionary analysis of ancient genome duplication in the yeast *Saccharomyces cerevisiae*. *Nature* 428:617–624.
29. Jaillon O, et al. (2004) Genome duplication in the teleost fish *Tetraodon nigroviridis* reveals the early vertebrate proto-karyotype. *Nature* 431:946–957.
30. The Arabidopsis Genome I (2000) Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* 408:796–815.
31. Dietrich FS, et al. (2004) The *Ashbya gossypii* genome as a tool for mapping the ancient *Saccharomyces cerevisiae* genome. *Science* 304:304–307.
32. Gladyshev E, Meselson M (2008) Extreme resistance of bdelloid rotifers to ionizing radiation. *Proc Natl Acad Sci USA* 105:5139–5144.
33. Cox MM, Battista JR (2005) *Deinococcus radiodurans*, the consummate survivor. *Nat Rev Microbiol* 3:882–892.
34. Pagani M, Ricci C, Redi CA (1993) Oogenesis in *Macrotrachela quadricornifera*. I. Germarium eutely, karyotype, and DNA content. *Hydrobiologia* 255/256:225–230.
35. Mark Welch DB, Meselson M (2003) Oocyte nuclear DNA content and GC proportion in rotifers of the anciently asexual class Bdelloidea. *Biol J Linn Soc London* 79:85–91.
36. Mark Welch DB, Meselson M (1998) Measurements of the genome size of the monogonont rotifer *Brachionus plicatilis* and of the bdelloid rotifers *Philodina roseola* and *Habrotrocha constricta*. *Hydrobiologia* 387/388:395–402.
37. Stephens JC (1985) Statistical methods of DNA sequence analysis: Detection of intra-genic recombination or gene conversion. *Mol Biol Evol* 2:539–556.
38. Kirkpatrick M, Jenkins CD (1989) Genetic segregation and the maintenance of sexual reproduction. *Nature* 339:300–301.
39. Chasnov JR (2000) Mutation-selection balance, dominance and the maintenance of sex. *Genetics* 156:1419–1425.
40. White MJD (1973) *Animal Cytology and Evolution* (Cambridge Univ Press, Cambridge, UK).
41. Lee S, Rasheed S (1990) A simple procedure for maximum yield of high-quality plasmid DNA. *BioTechniques* 9:676–679.
42. Eddy SR (1998) Profile hidden Markov models. *Bioinformatics* 14:755–763.
43. Birney E, Clamp M, Durbin R (2004) GeneWise and genomewise. *Genome Res* 14:988–995.
44. Pamilo P, Bianchi NO (1993) Evolution of the Zfx and Zfy genes: Rates and interdependence between the genes. *Mol Biol Evol* 10:271–281.